



**ANWAR ENT**  
The Right Choice For Your Health

## Mr. Mohiemen Anwar MBBS PhD DOHNS FRCS (ORL-HNS)

Consultant ENT Surgeon | Chelsea and Westminster Hospital | Phone: 0207 117 2437 | Mob: 07541 51 6454  
E-Mail: secretary@anwar-ent.com | The Westminster Wing outpatient, Chelsea and Westminster Private Care,  
Chelsea and Westminster NHS Foundation Trust SW10 9NH. GMC 6027494 FAX. 03300539680

### Gaviscon Advance



**UNLIKE ANTACIDS,  
IT WORKS IN TWO WAYS...**

**2. NEUTRALISING  
ACIDS FOR  
INDIGESTION  
RELIEF**



**1. RAFT  
PREVENTING  
REFLUX**



## WHAT IS GAVISCON ADVANCE?

Each 10 ml of aniseed flavoured Gaviscon Advance contains sodium alginate Ph Eur 1000 mg and potassium bicarbonate Ph Eur 200 mg as the active ingredients. The other ingredients are calcium carbonate, carbomer, methyl and propyl hydroxybenzoates, sodium saccharin, sodium hydroxide, aniseed flavour derived from fennel, and purified water.

Gaviscon Advance belongs to a group of medicines called "reflux suppressants". Reflux is a process in which the acid stomach juices flow back up into the food pipe.

Unlike the stomach, the lining of the food pipe is not resistant to acid, so that when reflux occurs

Heartburn and acid indigestion often occur during pregnancy or if you have a medical condition called "hiatus hernia".

Heartburn and acid indigestion can be made worse if you eat fatty or very spicy foods, drink too much alcohol, smoke, wear tight clothing, are overweight or when you bend forward or lie down.

## BEFORE TAKING GAVISCON ADVANCE

You can take Gaviscon Advance if you are pregnant or breastfeeding.

Each 10 ml dose contains 4.6 mmol of sodium, 2.0 mmol of potassium and 2.0 mmol of calcium. If you have been advised to follow a diet restricted in any of these please consult your doctor before taking this product.

This product contains methyl and propyl parabens which may rarely cause an allergic reaction.

This treatment may be less effective if you have decreased amounts of gastric acid produced in your stomach.

## HOW TO USE GAVISCON ADVANCE

Adults, including the elderly, and children 12 years and over: 5-10 ml (one to two 5 ml spoonfuls) after meals and at bedtime, or as directed.



Children under twelve years: Should only be taken on medical advice.

#### AFTER TAKING GAVISCON ADVANCE

If you forget to take a dose it is not necessary to double the dose next time. Just carry on taking the medicine at the recommended dose. If symptoms persist after seven days consult your doctor.

If you take too much Gaviscon Advance it is unlikely to cause you any harm. However, you may feel bloated. Consult your doctor or pharmacist if this feeling does not go away.

#### WHAT SIDE-EFFECTS MAY OCCUR?

Do not take if you know you are allergic to any of the ingredients as skin rashes and difficulty in breathing have been reported. Consult your doctor or pharmacist if you experience any side-effects after taking Gaviscon Advance.

#### STORAGE

Do not use this medicine after the expiry date (EXP: month/year) shown on this pack.

Keep all medicines safely away from children.

Do not refrigerate. Shake well before use.

Check that the cap seal is unbroken before first taking this product.